For Veterinary Use Only

E-SEVİL

Oral Emulsion

Veterinary Vitamin – Mineral

COMPOSITION

It is a white colored emulsion containing 1 mg sodium selenite, 60 mg Vitamin E and 40 mg Vitamin B1 per ml.

PHARMACOLOGICAL PROPERTIES

Selenium is one of the essential trace elements required by the body. It is necessary for normal growth and reproduction. It is used in the prevention and treatment of many degenerative diseases in domestic animals. Its effects are similar to vitamin E in many ways. Both substances complement each other's effects. Vitamin E has biological antioxidant properties. Vitamin B1 ensures normal development, regulates digestion and absorption, and acts as a coenzyme in carbohydrate metabolism. It has a preventive effect on functional disorders in the nervous system. Since certain amounts of selenium and vitamin E must be obtained externally along with the ration, their deficiencies are important for all domestic animals. E-sevil Oral Emulsion is a drug prepared for the prevention and treatment of disorders that may arise due to these deficiencies.

AREA OF USE/INDICATIONS

It is used for prevention and treatment of encephalomass, muscular dystrophy, and exudative diathesis in poultry and in reducing infertility rates caused by the deficiency or absence of active ingredients in the formulation in breeding flocks.

USAGE AND DOSAGE

It is administered orally in poultry. Shake the bottle before use.

2 ml product is administered by adding into 5 L water. If needed, the dose is repeated a week later.

In continuous treatment, maximum 4 doses can be administered at 2-week intervals.

SPECIAL CLINICAL INFORMATION AND SPECIAL WARNINGS FOR TARGET SPECIES

Some animals may be sensitive to selenium; a test dose should be administered prior to use. In sensitive animals, half the dose may be administered on two consecutive days. As selenium is a toxic substance, therapeutic doses should not be exceeded. Horses are sensitive to selenium.

ADVERSE EFFECTS

Hypersensitivity reactions are rarely seen in cattle. In such cases, symptomatic treatment is administered. Temporary irritation may occur at the injection site.

DRUG INTERACTIONS
There is no known incompatibility with the other drugs.

SYMPTOMS OF OVERDOSE, PRECAUTIONS AND ANTIDOTE
When used in high doses, increased respiratory rate, respiratory distress, tachycardia, diarrhea, behavioral disorders, and collapse may occur. Symptomatic treatment is administered in cases of overdose. There is no specific antidote.

WITHDRAWAL PERIODS
'0' day for meat and egg.

CONTRAINDICATIONS
There is no known contraindications with the other drugs.

GENERAL WARNINGS
Consult a veterinarian before use and if any unexpected reactions occur.
Keep out of reach of children and away from foodstuffs.

PRECAUTIONS TO BE TAKEN BY THE ADMINISTRATOR AND WARNINGS FOR VETERINARY SURGEON
Avoid contact with the drug and wash your hands after using.

STORAGE CONDITIONS AND SHELF LIFE
It should be stored in the original packages, protected from light, below 25°C, without refrigeration or freezing. Shelf life is 3 years.

COMMERCIAL PRESENTATION FORM
100 ml HDPE plastic bottles with boxes and 1 L white HDPE plastic bottles without boxes.

PLACE OF SALE AND CONDITIONS
It is sold in pharmacies, veterinary surgeries, polyclinics and animal hospitals with a veterinarian's prescription. (VSP)

APPROVAL DATE OF LEAFLET: 23.12.2015

AUTHORIZATION DATE AND NO: 23.12.2015 – 132

NAME AND ADDRESS OF AUTHORIZATION HOLDER
VİLSAN Veteriner İlaçları Tic. ve San. A.Ş.
Balıkhisar Mah. Balıkhisar Köy İçi Küme Evleri No:765A Akyurt/Ankara-Türkiye

MANUFACTURER COMPANY
VİLSAN Veteriner İlaçları Tic. ve San. A.Ş.
Balıkhisar Mah. Balıkhisar Köy İçi Küme Evleri No:765A Akyurt/Ankara-Türkiye

Made in Turkiye