

## **Leaflet**

For Veterinary Use Only  
**OPTIVIL**  
Eye Ointment  
Veterinary Topical Antibacterial

### **COMPOSITION**

It is a white-colored cream ointment containing 213 mg cloxacillin benzathine.

### **PHARMACOLOGICAL PROPERTIES**

It is a local and long-acting antibacterial. Due to the activity of cloxacillin benzathine against penicillinase, it is particularly effective against bacteria resistant to Penicillin G (penicillinase-producing *Staphylococci*). It exhibits bactericidal activity.

Susceptible bacteria include *Staphylococcus spp.* (including beta-lactamase-producing strains) and *Bacillus spp.*

After ocular administration, cloxacillin benzathine maintains a concentration above the minimum inhibitory concentration (MIC) in the lacrimal fluid for 2–3 days, providing effective levels in the eye. It is eliminated via tear fluid.

### **AREA OF USE/INDICATIONS**

It is used for the treatment of ocular infections locally in horses, cattle, sheep, cats and dogs.

### **USAGE AND DOSAGE**

**Cattle and horses:** 5–10 cm line is applied by opening the lower eyelid and squeezing it into the conjunctival sac for each eye.

**Sheep:** 5 cm strip for each eye.

**Cats and dogs:** 2 cm strip for each eye.

Normally, a single administration is sufficient. If necessary, treatment may be repeated within 48–72 hours.

To prevent cross-infection, it is recommended to also treat the non-infected eye. To minimize the risk of spreading infection, always begin treatment with the healthy eye.

For external use only.

### **ADVERSE EFFECTS**

No adverse effects have been reported at the recommended doses. Allergic reactions may occur in animals sensitive to penicillins.

### **DRUG INTERACTIONS**

Do not use concomitantly with tetracyclines.

### **WITHDRAWAL PERIODS**

**Meat and milk: zero (0) days.**

## **CONTRAINDICATIONS**

Do not use in animals known to be hypersensitive to penicillins or cephalosporins.

**Use during pregnancy:** It can be used safely during pregnancy and lactation.

## **GENERAL WARNINGS**

Consult a veterinarian before use and in case of unexpected reactions. Keep out of reach of children.

## **PRECAUTIONS TO BE TAKEN BY THE ADMINISTRATOR AND WARNINGS FOR VETERINARY SURGEON**

Avoid contact with the product and wash hands after use. People known to be hypersensitive to penicillins and cephalosporins should not handle this product. In sensitive people, exposure by oral, dermal, or inhalation routes may cause swelling of the face, eyes, or lips, or difficulty breathing; such symptoms require immediate medical attention. In such cases, contact a doctor and show the product label.

## **STORAGE CONDITIONS AND SHELF LIFE**

It should be stored in the original packages, protected from light, below 25°C, without refrigeration or freezing. Shelf life is 2 years.

## **COMMERCIAL PRESENTATION FORM**

5 g white plastic syringes, in box containing 24 units.

## **PLACE OF SALE AND CONDITIONS**

It is sold in pharmacies with a veterinary prescription, as well as in veterinary clinics, polyclinics and animal hospitals. (VSP)

## **APPROVAL DATE OF LEAFET: 24.04.2009**

## **MARKETING AUTHORIZATION DATE AND NO: 13.04.2009-021/0023**

## **MARKETING AUTHORIZATION HOLDER**

VİLSAN Veteriner İlaçları Tic. ve San. A.Ş.

Balıkhisar Mah. Balıkhisar Köy İçi Küme Evleri No:765A Akyurt/Ankara–Türkiye

## **MANUFACTURER COMPANY**

VİLSAN Veteriner İlaçları Tic. ve San. A.Ş.

Balıkhisar Mah. Balıkhisar Köy İçi Küme Evleri No:765A Akyurt/Ankara–Türkiye

**Made in Türkiye**

**Box**

For Veterinary Use Only  
**OPTİVİL**  
Eye Ointment  
Veterinary Topical Antibacterial

**COMPOSITION**

It contains 213 mg cloxacillin benzathine.

**AREA OF USE/INDICATIONS**

It is used for the treatment of ocular infections locally in horses, cattle, sheep, cats, and dogs.

**USAGE AND DOSAGE**

**Cattle and horses:** 5–10 cm line is applied by opening the lower eyelid and squeezing it into the conjunctival sac for each eye.

**Sheep:** 5 cm strip for each eye.

**Cats and dogs:** 2 cm strip for each eye.

Normally, a single administration is sufficient. If necessary, treatment may be repeated within 48–72 hours.

**WITHDRAWAL PERIODS**

**Meat and milk: zero (0) days.**

**GENERAL WARNINGS**

Consult a veterinarian before use and in case of unexpected reactions. Keep out of reach of children. Read the leaflet before use.

**STORAGE CONDITIONS AND SHELF LIFE**

It should be stored in the original packages, protected from light, below 25°C, without refrigeration or freezing.

**MARKETING AUTHORIZATION DATE AND NO: 13.04.2009-021/0023**

**MARKETING AUTHORIZATION HOLDER**

VİLSAN Veteriner İlaçları Tic. ve San. A.Ş.

Balıkhisar Mah. Balıkhisar Köy İçi Küme Evleri No:765A Akyurt/Ankara–Türkiye

**MANUFACTURER COMPANY**

VİLSAN Veteriner İlaçları Tic. ve San. A.Ş.

Balıkhisar Mah. Balıkhisar Köy İçi Küme Evleri No:765A Akyurt/Ankara–Türkiye

**Made in Türkiye**

**Label**

For Veterinary Use Only  
**OPTIVİL**  
Eye Ointment  
Veterinary Topical Antibacterial

It contains 213 mg cloxacillin benzathine.

It is used for the treatment of ocular infections locally in horses, cattle, sheep, cats and dogs.

**Cattle and horses:** 5–10 cm line is applied by opening the lower eyelid and squeezing it into the conjunctival sac for each eye. 5 cm strip for each eye for sheep, 2 cm strip for each eye for cat and dog.

**Meat and milk: zero (0) days.**

Consult a veterinarian before use and in case of unexpected reactions. Keep out of reach of children. Read the leaflet before use.

It should be stored in the original packages, protected from light, below 25°C, without refrigeration or freezing.

**MARKETING AUTHORIZATION HOLDER**

VİLSAN Veteriner İlaçları Tic. ve San. A.Ş.

Balıkhisar Mah. Balıkhisar Köy İçi Küme Evleri No:765A Akyurt/Ankara–Türkiye

**MANUFACTURER COMPANY**

VİLSAN Veteriner İlaçları Tic. ve San. A.Ş.

Balıkhisar Mah. Balıkhisar Köy İçi Küme Evleri No:765A Akyurt/Ankara–Türkiye

**Made in Türkiye**